



Achilles tendon rupture treated with VACOped® Boot: user guide

Information for patients

If you have any questions once you get home, please contact your Achilles clinical specialist physiotherapist on 01227 783065.

When removing your boot, you must keep your toes and ankle pointed down at all times. You may need someone to hold your foot to help you keep your ankle in this position.

Week 1 to 4: Boot fixed at 30°

You do not need to do anything for the first four weeks.



Boot fixed at 30°

Week 5 to 6: Range of movement 15° to 30°

How do I change it?

- Insert the VACOped® Key in the lower "keyhole".
- Turn the key 90° either way, and move the lower fixation three notches down from the previous setting.
- Lock it back by turning the key 90°. One notch = 5°.



Insert the VACOped® Key in the lower "keyhole"



Move the lower fixation three notches down from the previous setting

Week 7 to 8: Range of movement 0° to 30°, plus flat sole



Using the VACOped® Key, move the lower fixation three notches down from the previous setting



Remove the wedge sole, by pressing the "buttons" on both sides



Adjust the flat sole. You should be able to hear it "click" when in place

Other adjustments

If you feel the boot is too long or too short, you can adjust the length between the sole and the actual boot.



Adjusting the length between the sole and the actual boot

Looking after your boot

- You need to keep your boot dry.
- Make sure you use your pump every morning.
- No more than once a week, when removing your boot or changing the liner to wash it, you need to give the inner liner a good shake to spread the beads equally throughout the boot.

Changing and washing the cushion liner

- To change the cushion liner simply unzip it, remove the inner liner and put in a clean one.
- Make sure each section comes into the appropriate pouch in the cushion liner, and the valve comes out through the designated hole.
- Zip it back and put it on.

You can wash the cushion liner in the washing machine up to 60°C.

If your boot is creating pressure sores on your skin, you can place a dressing on the affected area(s).

If you feel your hips or back are suffering because of the uneven gait that wearing the boot causes, there are products that can help by providing extra elevation (height) under your non-affected limb.

For more information and explanatory videos, see the OPED UK (https://oped-uk.com/products/) web site.

This leaflet has been produced with and for patients.

If you would like this information in **another language, audio, Braille, Easy Read, or large print** please ask a member of staff. You can ask someone to contact us on your behalf.

Any complaints, comments, concerns, or compliments please speak to your doctor or nurse, or contact the Patient Advice and Liaison Service (PALS) on 01227 783145 (tel://+441227783145), or email ekh-tr.pals@nhs.net (ekh-tr.pals@nhs.net)

Patients should not bring large sums of money or valuables into hospital. Please note that East Kent Hospitals accepts no responsibility for the loss or damage to personal property, unless the property has been handed into Trust staff for safe-keeping.

Further patient leaflets are available via the East Kent Hospitals website (https://www.ekhuft.nhs.uk/information-for-patients/patient-information/).

Reference number: Web 477 (A4)

First published: July 2020 Last reviewed: January 2023 Next review date: May 2026

Copyright © East Kent Hospitals University NHS Foundation Trust.